

Endophytic Xylariaceae from Thai Plants: A Research Review

Anthony J S Whalley^{*1,5}, Nuttika Suwannasai², Nutthaporn Ruchikachorn³,
Ek Sangvichien⁴ and Prakitsin Sihanonth⁵

¹School of Pharmacy and Biomolecular Sciences,
Liverpool John Moores University, Liverpool, UK

²Department of Biology, Faculty of Science,
Srinakharinwirot University, Bangkok, Thailand

³State Key Laboratory of Mycology, Institute of Microbiology,
Chinese Academy of Sciences, Beijing 100101, P.R. China

⁴Department of Biology, Faculty of Science,
Ramkhamhaeng University, Bangkok, Thailand

⁵Department of Microbiology, Faculty of Science,
Chulalongkorn University, Bangkok, Thailand

*Corresponding Author: a.j.whalley@ljmu.ac.uk

Abstract: Endophytic fungi are present in all plant species investigated, and members of the Xylariaceae are especially common in tropical plants. There is major interest in endophytic fungi, because of their excellent track record concerning the production of novel and often bioactive compounds including the anti-cancer drugs taxol and taxane. Studies in Thailand on endophytes over the past 20 years confirm the high presence of the Xylariaceae in many different Thai plants. Species of *Xylaria* stand out as the most frequent Xylariaceae isolated; although, *Daldinia eschscholtzii* has also been commonly isolated. DNA technology and chemical profiling have been shown to be invaluable in the identification of xylariaceous isolates which previously could not be identified to species level or even assigned to genera in many cases. Reference is made to those isolates which produce novel compounds or those exhibiting bioactive properties.

Keywords: endophytic fungi, xylariaceae, Thailand, novel metabolites.

Introduction

Endophytic fungi have been described as those organisms that live inside the plant tissue for at least part of their life cycle without causing any disease symptoms in the host (Petrini, 1991). There is currently major interest in endophytes mainly following the discovery of the anticancer drugs taxol and taxane in the endophyte *Taxomyces andreanae* Strobel, A. Stierle, D. Stierle & W. M. Hess (Stierle *et al.*, 1993) and subsequent references to bioactive compounds isolated from endophytes (Strobel *et al.*, 2004; Tenguria *et al.*, 2011). It is noteworthy that the Xylariaceae are well known as a source of novel metabolites (Whalley & Edwards, 1995; Stadler & Hellwig, 2005) but also exhibit a remarkable presence as endophytes being regular and often dominant members of the endophytic communities of tropical plants (Petrini *et al.*, 1995). It is therefore not surprising that the Xylariaceae have been the target for many researchers. In Thailand, there have been a number of important studies with numerous publications on both novelty of

the chemicals produced and their bioactive properties. Early studies on xylariaceous endophytes encountered difficulties in identification, certainly to species level, since they fail to produce stable diagnostic features and most of the anamorphic endophytic isolates failed to develop their teleomorph in culture. In most cases, the cultures and anamorphs are insufficiently distinctive to enable confident identification but pioneering work by Petrini and colleagues (*e.g.* Petrini, 2013; Petrini & Petrini, 1985; Petrini *et al.*, 1995) has resulted in the development of keys and publication of suitable data to allow identification to be made at least to generic level for temperate isolates. The situation regarding tropical endophytic Xylariaceae is much more complex as a result of their abundance, almost universal presence and their impressive diversity (Rodrigues & Samuels, 1990; Rodrigues, 1994; Whalley, 1996). It is doubtful whether differentiation of species on the basis of cultural and anamorphic features alone will ever be possible since differences between individual species are often insufficient to allow

for absolute identifications to be made (Petrini *et al.*, 1995). Attempts were made to use a combination of morphological and biochemical data and these were considered to be promising (Brunner & Petrini, 1992; Rodrigues *et al.*, 1993). During this period, investigations into secondary metabolite profiles from endophytic isolates might be 'matched' with those obtained from cultures derived from teleomorphic material thus enabling identity to be established (Whalley & Edwards, 1995; Whalley, 1996). In a few cases, identification could be made on the basis of metabolite profiles compared with those from known teleomorph forms. Since then, Stadler and his many co-workers have developed a sophisticated database of secondary metabolites occurring in an impressive range of xylariaceous taxa using HPLC-DAD/MS profiling, HPLC-DAD/MS de-replication, GC-MS profiling and NMR spectroscopy (Stadler *et al.*, 2001a, 2001b, 2008, 2010). A recent monograph of the genus *Daldinia* Ces. & De Not. combining traditional taxonomic characteristics with chemical and molecular data provides powerful evidence for this approach and the great future potential for the identification of these previously unidentifiable xylariaceous endophytes (Stadler *et al.*, 2014).

In this mini review, we refer to important aspects of the lifestyle of endophytic fungi concentrating on our studies in Thailand over the past 20 years. Previously unpublished data is included and is discussed in the light of other studies on endophytic fungi.

Endophytic Xylariaceae of Thailand

In early studies of endophytes of *Tectona grandis* L (teak), attempts were made to produce the teleomorphs of a range of endophytic Xylariaceae to enable identifications to be made. It was found that inoculation of pre-sterilized twigs with pure cultures isolated from mature stromata on decaying plant material collected from the sites where endophytic isolates had been obtained or from cultures of selected xylariaceous endophytes which when placed in the forest to develop could often produce mature stromata (Mekkamol *et al.*, 1997; Mekkamol, 1998). *Daldinia eschscholtzii* (Ehrenb.) Rehm took 12-14 weeks, but species of *Xylaria* took over 24 weeks or longer to maturity. *Xylaria grammica* (Mont.) Mont., *X. cubensis* (Mont.) Fr. and *D. eschscholtzii* were subsequently recognized from mature stromata developing in twigs inoculated

with endophytic isolates. Interestingly *D. eschscholtzii* was found to be the dominant endophyte in teak leaves sampled early in the rainy season with *Xylaria* later becoming more frequent (Mekkamol *et al.*, 1997; Mekkamol, 1998). It is suggested that this was a result of the higher inoculum potential of the *Daldinia* ascospores following the development of mature stromata early in the rainy season and its ability to produce large numbers of spores. *Xylaria* species were found to take longer to develop to maturity and this might explain their low presence early in the rainy season but their greater frequency later on. Our more recent studies confirm that *D. eschscholtzii* is a frequent endophyte in Thai plants Chareprasert *et al.*, 2006, 2010, 2012). In their studies on endophytes of wild banana (*Musa acuminata*) at Doi Suthep Pui National Park (Photita *et al.*, 2001) and from *Amomum siamense* Xylariaceae were also common endophytic fungi (Bussaban *et al.*, 2001). A review of endophytic fungi in Thailand gave insight into their diversity and potential for future investigations (Lumyong *et al.*, 2004) Endophytes from *Garcinia* species also exhibited high xylariaceous representation (Phongpaichit *et al.*, 2006). A major study of endophytic Xylariaceae in Thailand was then undertaken by Okane *et al.* (2008) in which 405 strains of Xylariaceae (273 endophytic and 132 saprobic strains) were studied to examine the diversity and taxonomy of endophytes and to explore the relationships between those endophytes and saprobic Xylariaceae in Thailand recognized by their teleomorphic characteristics. In this study in KhaoYai National Park, it was found that *Xylaria* species were dominant and in analysis of 28S rDNA D1/D2 sequences 21 species of Xylariaceae were recorded inhabiting tropical plant foliage. Apart from *Xylaria*, *Hypoxylon haematostroma* Mont. and *D. eschscholtzii* were confirmed endophytic isolates. Our own studies from various sites and host plants in Thailand support these findings with *Xylaria* being by far the dominant xylariaceous genus in most cases (Mekkamol *et al.*, 1997; Chareprasert *et al.*, 2006, 2010, 2012; Pharamat *et al.*, 2013; Ruchikachorn, 2005).

Whalley (1996) reported *Anthostomella* Sacc., *Biscogniauxia* Kuntze, *Daldinia* Ces., *Hypoxylon* Bull., *Kretzschmaria* Fr., *Nemania* Gray, *Rosellinia* De Not. and *Xylaria* Hill ex Schrank to have endophytic representation. Members of these genera have been confirmed as endophytes in Thai plants and selected representatives are shown in Figure 1. Other genera since recorded

include *Annulohypoxyton* Y.-M. Ju, J.D. Rogers & H. M. Hsieh and *Muscodor* Worapong, Strobel & W. H. Hess. The genus *Muscodor albus* anam. gen. et sp. nov. was erected for an endophytic isolate from *Cinnamomum zeylandicum* and is included in the Xylariaceae on the basis of its molecular similarities (Worapong *et al.*, 2001). It appears to be widely distributed in tropical rainforests and has been recorded from Thailand (Sopalun *et al.*, 2003). The following new species of *Muscodor* have since been described from

Chiang Mai Province *M. cinnamomi* Suwannarach, Bussaban, K. D. Hyde & Lumyong, *M. musae* Suwannarach & Lumyong, *M. oryzae* Suwannarach & Lumyong, *M. suthepensis* Suwannarach & Lumyong and *M. equiseti* Suwannarach & Lumyong (Suwannarach *et al.*, 2010; 2013). *Muscodor* species are of considerable interest since they produce volatile metabolites with promise for biocontrol of spoilage fungi (Strobel, 2010).



Figure 1. Selected xylariaceous fungi. A. *Daldinia eschscholtzii*, B. *Kretzschmaria* sp., C. *Xylaria grammica*, D. *Xylaria cubensis*, E. *X. grammica* culture on PDA medium, F. *X. cubensis* culture on PDA medium. Bars = 1 cm.

Novel chemicals and bioactivity

Our early studies on secondary metabolites of the Xylariaceae demonstrated a remarkable diversity of chemicals with many proving to be novel structures (Adeboya *et al.*, 1995a, 1995b; Anderson *et al.*, 1982, 1983, 1984a, 1984b, 1985; Edwards & Whalley, 1979; Edwards *et al.*, 1988, 1989, 1991, 2001). It was also shown that the presence or absence of certain metabolites and metabolite profiles were of value in making taxonomic judgement (Whalley & Edwards, 1995). Since this early work, Stadler and co-workers have identified numerous new compounds and in many cases have demonstrated useful bioactive properties (Stadler *et al.*, 2001a, 2001b, 2008, 2010; Bitzer *et al.*, 2008). Their results have proven to be invaluable in both species identification and in taxonomic classification. These publications, together with the discovery of taxol and taxane in an endophyte, have resulted in wide ranging investigations of metabolites from the Xylariaceae and xylariaceous endophytes. There are too many publications to list all here, but a few selected papers on bio-active properties by Daferner *et al.*, 1999; Singh *et al.*, 1999; Isaka *et al.*, 2000; Quang *et al.*, 2006 provide a useful base to work from. Sodngam *et al.* (2014) investigated the chemicals and their activity from *Xylaria humosa* Lloyd recently found in Thailand and provide key references to bioactive compounds from *Xylaria* species. Stadler and Hellwig (2005) have reported on a wide range of chemicals identified and their bioactive properties.

Conclusions

The Xylariaceae are common and widely distributed endophytes in those Thai plants which have been investigated. The genus *Xylaria* is dominant in most of the studies undertaken but *D. eschscholtzii* is also a regular inhabitant and in certain situations can be numerically the most important. There are a large number of isolates recognized as xylariaceous without identification to genus and species levels. *Xylaria* endophytes are in particular a good source of metabolites, many novel and a number exhibit bioactive properties.

Acknowledgements

It is a pleasure to thank many colleagues and former students for their help and advice during our studies over the years.

References

- Adeboya, M., Edwards, R. L., Læssøe, T. & Maitland, D. J. (1995a). Metabolites of the higher fungi. Part 27. Berteric acid, cameronic acid and malaysic acid, three new polysubstituted fatty acids related to cubensic acid from species of the fungus genus *Xylaria*. *Journal of Chemical Research, Synopses*, 9, 356-357.
- Adeboya, M., Edwards, R. L., Læssøe, T., Maitland, D. J. & Whalley, A. J. S. (1995b). Metabolites of the higher fungi. Part 28. Globosic acid and globoscin, a labile acid-lactone system from *Xylaria globosa* and *Xylaria obovata*. *Journal of the Chemical Society. Perkins Transactions*, 1, 2067-2072.
- Anderson, J. R., Briant, C. E., Edwards, R. L., Mabelis, R. P., Poyser, J. P., Spencer, H. & Whalley, A. J. S. (1984a). Punctatin A (antibiotic M95464): X-ray crystal structure of a ses- quiterpene alcohol with a new carbon skeleton from the fungus *Poronia punctata*. *Journal of the Chemical Society, Chemical Communications*, 405-406.
- Anderson, J. R., Edwards, R. L., Freer, A. A., Mabelis, R. P., Poyser, J. P., Spencer, H. & Whalley, A. J. S. (1984b). Punctatins B and C (antibiotics M95154 and M951560): further sesquiterpene alcohols from the fungus *Poronia punctata*. *Journal of the Chemical Society, Chemical Communications*, 917-919.
- Anderson, J. R., Edwards, R. L. & Whalley, A. J. S. (1982). Metabolites of the higher fungi. Part 19. Serpenone, 3-methoxy-4-methyl-5 prop-1-enylfuran-2(5H)-one, a new γ - butyrolactone from the fungus *Hypoxylon serpens* (Barrons strain)(Persoon ex Fries)Kickx. *Journal of the Chemical Society, Perkins Transactions* 1, 215-221.
- Anderson, J. R., Edwards, R. L. & Whalley, A. J. S. (1983). Metabolites of the higher fungi. Part 21. 3-Methyl-3, 4-dihydroisocoumarins and related compounds from the ascomycete family Xylariaceae. *Journal of the Chemical Society. Perkins Transactions* 1, 2185-2192.
- Anderson, J. R., Edwards, R. L. & Whalley, A. J. S. (1985). Metabolites of the higher fungi. Part 22. 3-Butyl-3-methylsuccinic acid and a 2-hexylidene-3-methylsuccinic acid from xylariaceous fungi. *Journal of the Chemical Society. Perkins Transactions* 1, 1481-1485.
- Bitzer, J., Læssøe, T., Fournier, J., Kummer, V., Decock, C., Tichy, H. V., Piepenbring, M., Persoh, D. & Stadler, M. (2008). Affinities of *Phylacia* and the daldinoid Xylariaceae

- inferred from chemotypes of cultures and ribosomal DNA sequences. *Mycological Research*, 112 (2): 251-270
- Brunner, F. & Petrini, O. (1992). Taxonomy of some *Xylaria* species and xylariaceous endophytes by isozyme electrophoresis. *Mycological Research*, 96, 723-733.
- Bussaban, B., Lumyong, S., Lumyong, P., McKenzie, E. H. C. & Hyde, K. D. (2001). Endophytic fungi from *Amomum siamense*. *Canadian Journal of Microbiology*, 47, 943-948.
- Chareprasert, S., Abdelghany, M. T., El-sheikh, H. H., Ahmed, A. F., Khalil, A. M. A., Sharples, G. P., Sihanonth, P., Soliman, H. G., Suwannasai, N., Whalley, A. J. S. & Whalley, M. A. (2012). *Xylariaceae on the fringe*. In C. Raghukumar (Ed), *Biology of Marine Fungi* (pp. 229-241). Berlin, Heidelberg: Springer-Verlag Berlin Heidelberg.
- Chareprasert, S., Piapukiew, J., Thienhirun, S., Whalley, A. J. S. & Sihanonth, P. (2006). Endophytic fungi of teak leaves (*Tectona grandis* L. and rain tree leaves *Samanea saman* Merr. *World Journal of Biotechnology and Microbiology*, 22(5), 481-486.
- Chareprasert, S., Piapukiew, J., Whalley, A. J. S. & Sihanonth P. (2010). Endophytic fungi from selected mangrove plants in Chanthaburi, Prachuap Kiri Khan and Ranong provinces, their potential as a source of antimicrobial and anticancer compounds. *Botanica Marina*, 53, 555-564.
- Daferner, M., Mensch, S., Anke, T. & Sterner, O. (1999). Hypoxysordarin, a new sordarin derivative from *Hypoxylon croceum*. *Zeitschrift für Naturforschung C*, 54(7-8): 474-480.
- Edwards, R. L., Maitland, D. J., Scowen, I. A., De Sousa, A. J. T. & Whalley, A. J. S. (2001). Metabolites of the higher fungi. Part 32. Rosnecatrone, a phytotoxic Bicyclo [4.1.0] hept-3-en-2-one from the fungus *Rosellinianecatrix* Prill. *Journal of the Chemical Society, Perkins Transactions 1*, 537-542.
- Edwards, R. L., Maitland, D. J., Poyser, J. P. & Whalley, A. J. S. (1989). Metabolites of the higher fungi. Part 25. Puntaporonin G from the fungus *Poronia punctata* (Linneus: Fries) Fries. *Journal of the Chemical Society, Perkins Transactions 1*, 57-65.
- Edwards, R. L., Poyser, J. P. & Whalley, A. J. S. (1988). Metabolites of the higher fungi. Part 23. The puntaporonins. Novel bi-, tri, and tetra-cyclic sesquiterpenes related to caryophyllene from the fungus *Poronia punctata* (Linneus: Fries) Fries. *Journal of the Chemical Society, Perkins Transactions 1*, 823-831.
- Edwards, R. L. & Whalley, A. J. S. (1979). Metabolites of the higher fungi. Part 18. 3-butyl-4-methyl furan-2(5H)-one and 3-butyl-4-methylenefuran-2(5H)-one. New γ -butyrolactones from the fungus *Hypoxylon serpens* (Persoon ex Fries) Kickx. *Journal of the Chemical Society, Perkins Transactions 1*, 803-806.
- Edwards, R. L., Maitland, D. J. & Whalley, A. J. S. (1991). Metabolites of the higher fungi. Part 26. Cubensic acid, 3,7,11,15-tetrahydroxy-18-(hydromethyl)-2,4,6,10,16,20-heptamethyldocosa-4E,8E,12E,16E tetraenomic acid, a novel polysubstituted C22 fatty acid from the fungus *Xylaria cubensis* (Mont.) Fr. with substituent and substitution pattern similar to the macrolide antibiotics. *Journal of the Chemical Society, Perkins Transactions 1*, 1411-1417.
- Isaka, M., Jaturapat, A., Kladwang, W., Punya, J., Lertwerawat, Y., Tanticharoen, N. & Thebtaranonth, Y. (2000). Antiplasmodial compounds from the wood-decayed fungus *Xylaria* sp. BCC 1067. *Planta Medica*, 66(5), 473-475.
- Lumyong, S., Lumyong, P. & Hyde, K. D. (2004). *Endophytes*. In E. B. G. Jones, M. Tantichareon & K. D. Hyde (Eds.), *Thai fungal Diversity* (pp. 7-35). Bangkok: BIOTEC.
- Mekkamol, S. (1998). *Endophytic fungi in Tectona grandis* L. (Ph.D. Thesis). Liverpool, UK: Liverpool John Moores University.
- Mekkamol, S., Jonglaeka, N., Hywel-Jones, N. L. & Whalley, A. J. S. (1997). Endophytic fungi of teak in Thailand: a preliminary account. *BIOTEC News*, 3, pp. 4-5.
- Okane, I., Srikitikul, P., Toyama, K., Læssøe, T., Sivichai, S., Hywel-Jones, N., Nakagiri, A., Potacharoen, W. & Suzuki, K. L. (2008). Studies of endophytic Xylariaceae in Thailand: diversity and taxonomy inferred from rDNA sequence analyses with saprobes forming fruit bodies in the field. *Mycoscience*, 49, 359-372.
- Petrini L. E. & Petrini, O. (1985). Xylariaceous fungi as endophytes. *Sydowia*, 38, 216-234.
- Petrini, O. (1991). *Fungal endophytes of tree leaves*. In: *Microbial Ecology of Leaves* (eds Andrews, J. H. & Hirano, S. S.). : New York. Springer-Verlag.
- Petrini, O., Petrini, L. E. & Rodrigues, K. F. (1995). Xylariaceous endophytes: an exercise in bio-

- diversity. *Fitopatologia Brasileira*, 20, 531-539.
- Petrini, L. E. (2013). *Rosellinia—a world monograph*. Stuttgart: J. Cramer.
- Pharamat, T., Palaga, T., Piapukiew, J., Whalley, A. J. S. & Sihanonth, P. (2013). Antimicrobial and anticancer activities of endophytic fungi from *Mitrajyna javanica* Koord and Val. *African Journal of Microbiological Research*, 7, 5565-5572.
- Photita, W., Lumyong, S., Lumyong, P. & Hyde, K. D. (2001). Endophytic fungi of wild banana (*Musa acuminata*) at Doi Suthep Pui National Park, Thailand. *Mycological Research*, 105(12), 1508-1513.
- Phongpaichit, S., Rungjindamai, N., Rukachaisirikul V. & Sakayaroj, J. (2006). Antimicrobial activity in cultures of endophytic fungi isolated from *Garcinia* species. *FEMS Immunology and Medical Microbiology*, 48, 367-372.
- Quang, D. N., Stadler, M., Fournier J., Tomita, A. & Hashimoto, T. (2006). Cohaeirins C-F, four azaphilones from the xylariaceous fungus *Annulohypoxylon cohaerens*. *Tetrahedron*, 62, 6349-6354.
- Rodrigues, K. F. (1994). The foliar fungal endophytes of the Amazonian palm *Euterpe oleracea*. *Mycologia*, 86, 376-385.
- Rodrigues, K. F. & Samuels, G. J. (1990). Preliminary study of endophytic fungi in a tropical palm. *Mycological Research*, 94, 827-830.
- Rodrigues, K. F., Leuchtmann, A. & Petrini, O. (1993). Endophytic species of *Xylaria*: cultural and isozymic studies. *Sydowia*, 45, 116-138.
- Ruchikachorn, N. (2005). *Endophytic fungi of Cassia fistula L.* (Ph. D. Thesis). Liverpool, UK: Liverpool John Moores University.
- Singh, S. B., D. Zink, J. Polishook, D. Valentino & Shaffee, A. (1999). Structure and absolute stereochemistry of HIV-1 intergrase inhibitor integrin acid. A novel eremophilanesesquiterpenoid produced by a *Xylaria* sp. *Tetrahedron Letters*, 40(50), 8775-8779.
- Sodngam, S., Sawadsitang, S., Suwannasai, N. & Mongkolthanaruk, W. (2014). Chemical constituents, and their cytotoxicity, of the rare wood decay fungus *Xylaria humosa*. *Natural Products Chemistry*, 9, 157-158.
- Sopalun, M., Strobel, G. A., Hess, W. M. & Worapong, J. (2003). A record of *Muscodor albus*, an endophyte from *Myristica fragrans* in Thailand. *Mycotaxon*, 88, 239-247.
- Stadler, M. & Hellwig, V. (2005). Chemotaxonomy of the Xylariaceae and remarkable bioactive compounds from Xylariales and their associated asexual stages. *Recent Research Developments in Phytochemistry*, 9, 1-120.
- Stadler, M., Fournier, J., Gardt, S. & Persoh, D. (2010). The phylogenetic position of *Rhopalostroma* as inferred from a polythetic approach. *Persoonia*, 25, 11-21.
- Stadler, M., Fournier, J., Laessøe, L., Lechat, C., Tichy, H. & Piepenbring, M. (2008). Recognition of hypoxylid and xylarioid *Entonaema* species and allied *Xylaria* species from a comparison of holomorphic morphology, HPLC profiles, and ribosomal DNA sequences. *Mycological Progress*, 7, 53-73.
- Stadler, M., Læssøe, T., Fournier, J., Decock, C., Schmieschek, B., Tichy, H. V. & Persoh, D. (2014). A polyphasic taxonomy of *Daldinia* (Xylariaceae). *Studies in Mycology*, 77, 1-143.
- Stadler, M., Wollweber, H., Muhlbauer, A., Henke, T., Asakawa, Y., Hashimoto, T., Rogers J. D., Ju, Y. M., Wetzstein, H. G. & Tichy, H. V. (2001a). Molecular chemotaxonomy of *Daldinia* and other Xylariaceae. *Mycological Research*, 105, 1190-1205.
- Stadler, M., Wollweber, H., Muhlbauer, A., Henkel, T., Asakawa, Y., Hashimoto, T., Rogers, J. D., Ju, Y.-M., Wetzstein, H. G. & Tichy, H. V. (2001b). Secondary profiles, genetic fingerprints and taxonomy of *Daldinia* and allies. *Mycotaxon*, 77, 379-429.
- Stierle, A., Strobel, G. A. & Stierle, D. (1993). Taxol and taxane production by *Taxomyces andreanae*. *Science*, 260, 214-216.
- Strobel, G. (2010). *Muscodor* species—endophytes with biological promise. *Phytochemistry Reviews*, 10, 165-172.
- Strobel, G. A., Daisy, B., Castillo, U. & Harper, J. (2004). Natural products from endophytic microorganisms. *Journal of Natural Products*, 67, 257-268.
- Suannarach, N., Bussaban, B., Hyde, K. D. & Lumyong, S. (2010). *Muscodorc innamomi*, a new endophytic species from *Cinnamomum bejolghota*. *Mycotaxon*, 114, 14-23.
- Suannarach, N., Kumla, J., Bussaban, B., Hyde, K. D., Matsui K. & Lumyong S. (2013). Molecular evidence to support four new species in the genus *Muscodor* from northern Thailand. *Annals of Microbiology*, 63, 1341-1351.

- Tenguria, R. K., Khan, F. N. & Quereshi, S. (2011). Endophytesmines of pharmacological therapeutics. *World Journal of Science and Technology*, 1, 127-149.
- Whalley, A. J. S. (1996). The xylariaceous way of life. *Mycological Research*, 100, 897-922.
- Whalley, A. J. S. & Edwards, R. L. (1995). Secondary metabolites and systematic arrangement within the Xylariaceae. *Canadian Journal of Botany*, 73(Suppl. 1), S802-S810.
- Worapong, J., Strobel, G., Ford, E. J., Li, J. Y., Baird, G. & Hess, W. H. (2001). *Muscodor albus* anam. gen. et sp. nov. an endophyte from *Cinnamomum zeylandicum*. *Mycotaxon*, 79, 67-79.